



ALMA MATER STUDIORUM  
UNIVERSITÀ DI BOLOGNA

DIPARTIMENTO  
DI FARMACIA  
E BIOTECNOLOGIE

Progetto di Ricerca

**“Outcomes of early life experiences on neuroinflammatory-related processes in male and female adolescent rats”.**

Early social and emotional experiences differently shape individual's neurodevelopment inducing substantial changes in neurobiological substrates and behavior (Rullo et al., 2023).

Exposure to early life adversity (ELA) affects brain development differentially, depending on the type of adversity and the timing of exposure. ELA may trigger chronic neuroinflammation, which in turn affects key brain structures and functions, particularly in regions like hippocampus, amygdala, and prefrontal cortex (Andersen, 2022). In a similar way, early life stress (ELS) may induce long-lasting epigenetic modifications (e.g., DNA methylation) that increase vulnerability to neuroinflammation later in life, perpetuating a cycle of stress and immune dysregulation. For instance, it has been shown that overactivation of the amygdala due to neuroinflammation can lead to heightened emotional reactivity and increased susceptibility to anxiety disorders. Chronic stress and neuroinflammation may also impair the development of the prefrontal cortex, which is associated with decision-making, impulse control, and attention. In this frame, epigenetic mechanisms, including the activity of Jmjd3 (or KDM6A, histone demethylase), play a pivotal role in how early-life experiences like maternal separation have long-lasting effects on brain development. Maternal separation (an early-life stressor) likely triggers the activation of Jmjd3 in brain regions like the prefrontal cortex and hippocampus. Once activated, Jmjd3 demethylates repressive histone marks (H3K27me3), leading to the expression of genes involved in inflammatory pathways. This includes cytokines and other inflammatory mediators. Therefore, early-life stress can "program" certain genes to become more or less active later in life through epigenetic modifications. For example, Jmjd3's activation may be an adaptive response to stress in early life, but over time, its role in promoting neuroinflammation may become maladaptive, contributing to vulnerability to psychiatric disorders (Wang et al., 2020).

Neuroinflammation also influences opioid and cannabinoid receptor signaling, as chronic inflammation can desensitize or downregulate these receptors, contributing to the dysregulation of pain and reward circuits.

**SETTORE/UFFICIO/SEGRETERIA/DIREZIONE/LOREM IPSUM**

Via Massarenti 00 | 40100 Bologna | Italia | Tel. + 39 051 0000000 | email.@unibo.it



ALMA MATER STUDIORUM  
UNIVERSITÀ DI BOLOGNA

DIPARTIMENTO  
DI FARMACIA  
E BIOTECNOLOGIE

The endocannabinoid and opioid systems play protective roles in regulating stress and neuroinflammation, and their dysregulation due to epigenetic changes from ELS can have long-term consequences. The combined effects of epigenetic changes, neuroinflammation, and dysregulation of the endocannabinoid and opioid systems create a feedback loop that increases vulnerability to various psychiatric and neurological disorders (Catale et al., 2022).

It has previously been demonstrated that KDM6A increases in response to proinflammatory signaling, influencing the transcription of genes that PPARs regulate. PPAR $\alpha$  and PPAR $\gamma$ , important in reducing inflammation and metabolic stress, are modulated by the activity of KDM6A. When KDM6A is inhibited, there is an increase in the expression of PPARs, indicating a balancing act between these molecules to control neuroinflammatory processes (Rullo et al., 2021).

Based on this evidence, the aim of this research project will be the evaluation of potential molecular alteration induced by early life stress on KDMs, PPARs, and neuroinflammation mediators such as ILs, TNF- $\alpha$ , NLRP3, as well as endogenous opioids and their receptors. To assess this aim, molecular analysis will be performed in stress- and emotional-related brain area (hippocampus, amygdala, and prefrontal cortex) of adolescent male and female rats housed in either standard condition or subjected to a protocol of early social isolation (ESI, 30 min/day, during post-natal day 14-21).

The results obtained from this study will help to better understand the mechanisms of neuroinflammatory processes associated with early life experience and their impact on neurodevelopment.

## References

Andersen SL. Neuroinflammation, Early-Life Adversity, and Brain Development. *Harv Rev Psychiatry*. 2022 Jan-Feb 01;30(1):24-39. doi: 10.1097/HRP.0000000000000325. PMID: 34995033; PMCID: PMC8820591.

Catale C, Carola V, Viscomi MT. Early life stress-induced neuroinflammation and neurological disorders: a novel perspective for research. *Neural Regen Res*. 2022 Sep;17(9):1971-1972. doi: 10.4103/1673-5374.335152. PMID: 35142680; PMCID: PMC8848617.



ALMA MATER STUDIORUM  
UNIVERSITÀ DI BOLOGNA

DIPARTIMENTO  
DI FARMACIA  
E BIOTECNOLOGIE

Rullo L, Franchi S, Amodeo G, Caputi FF, Verduci B, Losapio LM, Sacerdote P, Romualdi P, Candeletti S. Interplay between Prokineticins and Histone Demethylase KDM6A in a Murine Model of Bortezomib-Induced Neuropathy. *Int J Mol Sci.* 2021 Nov 3;22(21):11913. doi: 10.3390/ijms222111913. PMID: 34769347; PMCID: PMC8584499.

Rullo L, Losapio LM, Morosini C, Mottarlini F, Schiavi S, Buzzelli V, Ascone F, Ciccocioppo R, Fattore L, Caffino L, Fumagalli F, Romualdi P, Trezza V, Candeletti S. Outcomes of early social experiences on glucocorticoid and endocannabinoid systems in the prefrontal cortex of male and female adolescent rats. *Front Cell Neurosci.* 2023 Dec 20;17:1270195. doi:

10.3389/fncel.2023.1270195. PMID: 38174157; PMCID: PMC10762649.

Wang R, Wang W, Xu J, Liu D, Wu H, Qin X, Jiang H, Pan F. Jmjd3 is involved in the susceptibility to depression induced by maternal separation via enhancing the neuroinflammation in the prefrontal cortex and hippocampus of male rats. *Exp Neurol.* 2020 Jun;328:113254. doi: 10.1016/j.expneurol.2020.113254. Epub 2020 Feb 19. PMID: 32084453.